

Black hole entropy and quantum information ¹

M. J. Duff ^{2†} and S. Ferrara^{3‡}

[†] *The Blackett Laboratory, Imperial College London,
 Prince Consort Road, London SW7 2AZ
 and*

[‡]*Physics Department, Theory Unit, CERN, CH1211, Geneva23, Switzerland and
 INFN-Laboratori Nazionale di Frascati, Via E. Fermi 40, 00044 Frascati, Italy*

ABSTRACT

We review some recently established connections between the mathematics of black hole entropy in string theory and that of multipartite entanglement in quantum information theory. In the case of $N = 2$ black holes and the entanglement of three qubits, the quartic $[SL(2)]^3$ invariant, Cayley's hyperdeterminant, provides both the black hole entropy and the measure of tripartite entanglement. In the case of $N = 8$ black holes and the entanglement of seven qubits, the quartic E_7 invariant of Cartan provides both the black hole entropy and the measure of a particular tripartite entanglement encoded in the Fano plane.

¹To appear in the proceedings of the “2006 Winter School on Attractor Mechanisms” (Frascati, Italy, March 2006), to be published in “Lecture Notes in Physics”, Springer.

²m.duff@imperial.ac.uk

³Sergio.Ferrara@cern.ch

Contents

1	Black holes and qubits	3
2	The N=2 STU model	4
3	Cayley's hyperdeterminant	7
4	Black hole entropy	8
5	The N=8 generalization	9
6	Decomposition of $E_{7(7)}$	10
7	Tripartite entanglement of 7 qubits	12
8	Cartan's $E_{7(7)}$ invariant	13
9	The black hole analogy	17
10	Subsectors	19
11	Conclusions	20
12	Acknowledgement	21

1 Black holes and qubits

It sometimes happens that two very different areas of theoretical physics share the same mathematics. This may eventually lead to the realisation that they are, in fact, dual descriptions of the same physical phenomena, or it may not. Either way, it frequently leads to new insights in both areas. In this paper the two areas in question are black hole entropy in string theory and qubit entanglement in quantum information theory. Going one way, we shall learn that the entropy of the so-called STU $N = 2$ black hole is given by the “hyperdeterminant”, a quantity first introduced by Cayley in 1845 and which describes the tripartite entanglement of three qubits [1, 2, 3]. Going the other way, we discover that the exceptional group E_7 , the U-duality group of $N = 8$ supergravity, plays a part in the tripartite entanglement of seven qubits [4, 5].

We begin in section 2 with an interesting subsector of string compactification to four dimensions which is provided by the STU model whose low energy limit is described by $N = 2$ supergravity coupled to three vector multiplets. One may regard it as a truncation of an $N = 4$ theory obtained by compactifying the heterotic string on T^6 where S, T, U correspond to the dilaton/axion, complex Kahler form and complex structure fields respectively. It exhibits an $SL(2, Z)_S$ strong/weak coupling duality and an $SL(2, Z)_T \times SL(2, Z)_U$ target space duality. By string/string duality, this is equivalent to a Type IIA string on $K3 \times T^2$ with S and T exchanging roles [6, 7, 8]. Moreover, by mirror symmetry this is in turn equivalent to a Type IIB string on the mirror manifold with T and U exchanging roles. Another way to obtain this model is by truncation of the $N=8$ theory that results from T^7 compactification of M-theory. Either way, the truncated theory has a combined $[SL(2, Z)]^3$ duality and complete $S - T - U$ triality symmetry [9]. Alternatively, one may simply start with this $N = 2$ theory directly as an interesting four-dimensional supergravity in its own right, as described in section 2.

The model admits extremal black holes solutions carrying four electric and magnetic charges and we organize these 8 charges into the $2 \times 2 \times 2$ *hypermatrix*, a_{ABD} , and display the $S - T - U$ symmetric Bogomolnyi mass formula [9]. Associated with this hypermatrix is a *hyperdeterminant*, $\text{Det } a_{ABD}$, discussed in section 3, first introduced by Cayley in 1845 [10]. The black hole entropy, first calculated in [11], is quartic in the charges and must be invariant under $[SL(2, Z)]^3$ and under triality. The main result of section 4, is to show [1] that this entropy given by the square root of Cayley’s hyperdeterminant:

$$S = \pi \sqrt{|\text{Det } a_{ABD}|}. \quad (1.1)$$

The hyperdeterminant also makes its appearance in quantum information theory [13]. Let the three qubit system ABD (Alice, Bob and Daisy) be in a pure state $|\Psi\rangle$, and let the components of $|\Psi\rangle$ in the standard basis be a_{ABD} :

$$|\Psi\rangle = a_{ABD} |ABD\rangle \quad (1.2)$$

or

$$\begin{aligned} |\Psi\rangle = & a_{000}|000\rangle + a_{001}|001\rangle + a_{010}|010\rangle + a_{011}|011\rangle \\ & + a_{100}|100\rangle + a_{101}|101\rangle + a_{110}|110\rangle + a_{111}|111\rangle \end{aligned} \quad (1.3)$$

Then the three way entanglement of the three qubits A , B and D is given by the *3-tangle* [12]

$$\tau_3(ABD) = 4|\text{Det } a_{ABD}|. \quad (1.4)$$

The 3-tangle is maximal for the GHZ state $|000\rangle + |111\rangle$ [28] and vanishes for the states $p|100\rangle + q|010\rangle + r|001\rangle$. The relation between three qubit quantum entanglement and the Cayley hyperdeterminant was pointed out by Miyake and Wadati [13].

As far as we can tell [1], the appearance of the Cayley hyperdeterminant in these two different contexts of stringy black hole entropy (where the a_{ABD} are integers and the symmetry is $[SL(2, Z)]^3$) and three-qubit quantum entanglement (where the a_{ABD} are complex numbers and the symmetry is $[SL(2, C)]^3$) is a purely mathematical coincidence. Nevertheless, it has already provided fascinating new insights [1, 2, 3, 4, 5] into the connections between strings, black holes, and quantum information⁴.

In section 6 we extend the argument to the $N = 8$ case and, noting that

$$E_{7(7)}(Z) \supset [SL(2, Z)]^7 \quad (1.5)$$

and

$$E_7(C) \supset [SL(2, C)]^7, \quad (1.6)$$

show that the corresponding system in quantum information theory is that of seven qubits (Alice, Bob, Charlie, Daisy, Emma, Fred and George). However, the larger symmetry requires that they undergo at most tripartite entanglement of a very specific kind. As discussed in section 8, the entanglement measure will be given by the quartic Cartan $E_7(C)$ invariant [16, 17, 18, 19]. The entanglement may be represented by the Fano plane [15] which also provides the multiplication table of the octonions. See also the interesting paper by Levay [5] who noted independently the connection to the Fano plane.

2 The N=2 STU model

Consider the three complex scalars axion/dilaton field S , the complex Kahler form field T and the complex structure field U

$$\begin{aligned} S &= S_1 + iS_2 \\ T &= T_1 + iT_2 \\ U &= U_1 + iU_2. \end{aligned} \quad (2.1)$$

This complex parameterization allows for a natural transformation under the various $SL(2, Z)$ symmetries. The action of $SL(2, Z)_S$ is given by

$$S \rightarrow \frac{aS + b}{cS + d}, \quad (2.2)$$

⁴A third application [14], not considered in this paper, is the Nambu-Goto string whose action is also given by $\sqrt{|\text{Det } a_{ABD}|}$ in spacetime signature (2, 2).

where a, b, c, d are integers satisfying $ad - bc = 1$, with similar expressions for $SL(2, Z)_T$ and $SL(2, Z)_U$. Defining the matrices $\mathcal{M}_S$, $\mathcal{M}_T$ and $\mathcal{M}_U$ via

$$\mathcal{M}_S = \frac{1}{S_2} \begin{pmatrix} 1 & S_1 \\ S_1 & |S|^2 \end{pmatrix}, \quad (2.3)$$

the action of $SL(2, Z)_S$ now takes the form

$$\mathcal{M}_S \rightarrow \omega_S^T \mathcal{M}_S \omega_S, \quad (2.4)$$

where

$$\omega_S = \begin{pmatrix} d & b \\ c & a \end{pmatrix}, \quad (2.5)$$

with similar expressions for $\mathcal{M}_T$ and $\mathcal{M}_U$. We also define the $SL(2, Z)$ invariant tensors

$$\epsilon_S = \epsilon_T = \epsilon_U = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (2.6)$$

Starting from the heterotic string, the bosonic action for the graviton $g_{\mu\nu}$, dilaton η , two-form $B_{\mu\nu}$ four $U(1)$ gauge fields A_S^a and two complex scalars T and U is [9]

$$\begin{aligned} I_{STU} = & \frac{1}{16\pi G} \int d^4x \sqrt{-g} e^{-\eta} \left[R_g + g^{\mu\nu} \partial_\mu \eta \partial_\nu \eta - \frac{1}{12} g^{\mu\lambda} g^{\nu\tau} g^{\rho\sigma} H_{\mu\nu\rho} H_{\lambda\tau\sigma} \right. \\ & + \frac{1}{4} \text{Tr}(\partial \mathcal{M}_T^{-1} \partial \mathcal{M}_T) + \frac{1}{4} \text{Tr}(\partial \mathcal{M}_U^{-1} \partial \mathcal{M}_U) \\ & \left. - \frac{1}{4} F_{S\mu\nu}^T (\mathcal{M}_T \times \mathcal{M}_U) F_S^{\mu\nu} \right]. \end{aligned} \quad (2.7)$$

where the metric $g_{\mu\nu}$ is related to the four-dimensional canonical Einstein metric $g_{\mu\nu}^c$ by $g_{\mu\nu} = e^\eta g_{\mu\nu}^c$ and where

$$H_{\mu\nu\rho} = 3(\partial_{[\mu} B_{\nu\rho]} - \frac{1}{2} A_{S[\mu}^T (\epsilon_T \times \epsilon_U) F_{S\nu\rho]}). \quad (2.8)$$

This action is manifestly invariant under T -duality and U -duality, with

$$F_{S\mu\nu} \rightarrow (\omega_T^{-1} \times \omega_U^{-1}) F_{S\mu\nu}, \quad \mathcal{M}_{T/U} \rightarrow \omega_{T/U}^T \mathcal{M}_{T/U} \omega_{T/U}, \quad (2.9)$$

and with η , $g_{\mu\nu}$ and $B_{\mu\nu}$ inert. Its equations of motion and Bianchi identities (but not the action itself) are also invariant under S -duality (2.2), with T and $g_{\mu\nu}^c$ inert and with

$$\begin{pmatrix} F_{S\mu\nu}^a \\ \tilde{F}_{S\mu\nu}^a \end{pmatrix} \rightarrow \omega_S^{-1} \begin{pmatrix} F_{S\mu\nu}^a \\ \tilde{F}_{S\mu\nu}^a \end{pmatrix}, \quad (2.10)$$

where

$$\tilde{F}_{S\mu\nu}^a = -S_2 [(\mathcal{M}_T^{-1} \times \mathcal{M}_U^{-1}) (\epsilon_T \times \epsilon_U)]^a_b * F_{S\mu\nu}^b - S_1 F_{S\mu\nu}^a, \quad (2.11)$$

where the axion field a is defined by

$$\epsilon^{\mu\nu\rho\sigma} \partial_\sigma a = \sqrt{-g} e^{-\eta} g^{\mu\sigma} g^{\nu\lambda} g^{\rho\tau} H_{\sigma\lambda\tau}, \quad (2.12)$$

and where $S = S_1 + iS_2 = a + ie^{-\eta}$.

Thus T -duality transforms Kaluza-Klein electric charges (F_S^3, F_S^4) into winding electric charges (F_S^1, F_S^2) (and Kaluza-Klein magnetic charges into winding magnetic charges), U -duality transforms the Kaluza-Klein and winding electric charge of one circle (F_S^3, F_S^2) into those of the other (F_S^4, F_S^1) (and similarly for the magnetic charges) but S -duality transforms Kaluza-Klein electric charge (F_S^3, F_S^4) into winding magnetic charge $(\tilde{F}_S^3, \tilde{F}_S^4)$ (and winding electric charge into Kaluza-Klein magnetic charge). In summary we have $SL(2, Z)_T \times SL(2, Z)_U$ and $T \leftrightarrow U$ off-shell but $SL(2, Z)_S \times SL(2, Z)_T \times SL(2, Z)_U$ and an S - T - U interchange on-shell.

One may also consider the Type IIA action I_{TUS} and the Type IIB action I_{UST} obtained by cyclic permutation of the fields S, T, U . Finally, one may consider an action [11] where the S, T and U fields enter democratically with a prepotential

$$F = STU \quad (2.13)$$

which off-shell has the full STU interchange but none of the $SL(2, Z)$. All four versions are on-shell equivalent.

Following [9], it is now straightforward to write down an S - T - U symmetric Bogomolnyi mass formula. Let us define electric and magnetic charge vectors α_S^a and β_S^a associated with the field strengths F_S^a and $\tilde{F}_S^a$ in the standard way. The electric and magnetic charges Q_S^a and P_S^a are given by

$$F_{S0r}^a \sim \frac{Q_S^a}{r^2} \quad * F_{S0r}^a \sim \frac{P_S^a}{r^2}, \quad (2.14)$$

giving rise to the charge vectors

$$\begin{pmatrix} \alpha_S^a \\ \beta_S^a \end{pmatrix} = \begin{pmatrix} S_2^{(0)} \mathcal{M}_T^{-1} \times \mathcal{M}_U^{-1} & S_1^{(0)} \epsilon_T \times \epsilon_U \\ 0 & -\epsilon_T \times \epsilon_U \end{pmatrix}^{ab} \begin{pmatrix} Q_S^b \\ P_S^b \end{pmatrix}. \quad (2.15)$$

For our purpose it is useful to define a $2 \times 2 \times 2$ array a_{ABD} via

$$\begin{pmatrix} a_{000} \\ a_{001} \\ a_{010} \\ a_{011} \\ a_{100} \\ a_{101} \\ a_{110} \\ a_{111} \end{pmatrix} = \begin{pmatrix} -\beta_S^1 \\ -\beta_S^2 \\ -\beta_S^3 \\ -\beta_S^4 \\ \alpha_S^1 \\ \alpha_S^2 \\ \alpha_S^3 \\ \alpha_S^4 \end{pmatrix}, \quad (2.16)$$

transforming as

$$a^{ABD} \rightarrow \omega_S^A \omega_T^B \omega_U^D a^{A'B'D'}. \quad (2.17)$$

Then the mass formula is

$$m^2 = \frac{1}{16} a^T (\mathcal{M}_S^{-1} \mathcal{M}_T^{-1} \mathcal{M}_U^{-1} - \mathcal{M}_S^{-1} \epsilon_T \epsilon_U - \epsilon_S \mathcal{M}_T^{-1} \epsilon_U - \epsilon_S \epsilon_T \mathcal{M}_U^{-1}) a. \quad (2.18)$$

This is consistent with the general $N = 2$ Bogomolnyi formula [34]. Although all theories have the same mass spectrum, there is clearly a difference of interpretation with electrically charged elementary states in one picture being solitonic monopole or dyon states in the other.

This $2 \times 2 \times 2$ array a_{ABD} is an example a “hypermatrix”, a term coined by Cayley in 1845 [10] where he also introduced a “hyperdeterminant”.

3 Cayley’s hyperdeterminant

In 1845 Cayley [10] generalized the determinant of a 2×2 matrix a_{AB} to the *hyperdeterminant* of a $2 \times 2 \times 2$ *hypermatrix* a_{ABD}

$$\begin{aligned} \text{Det } a &= -\frac{1}{2} \epsilon^{A_1 A_2} \epsilon^{B_1 B_2} \epsilon^{D_1 D_4} \epsilon^{A_3 A_4} \epsilon^{B_3 B_4} \epsilon^{D_2 D_3} a_{A_1 B_1 D_1} a_{A_2 B_2 D_2} a_{A_3 B_3 D_3} a_{A_4 B_4 D_4} \\ &= a_{000}^2 a_{111}^2 + a_{001}^2 a_{110}^2 + a_{010}^2 a_{101}^2 + a_{100}^2 a_{011}^2 \\ &\quad - 2(a_{000} a_{001} a_{110} a_{111} + a_{000} a_{010} a_{101} a_{111} \\ &\quad + a_{000} a_{100} a_{011} a_{111} + a_{001} a_{010} a_{101} a_{110} \\ &\quad + a_{001} a_{100} a_{011} a_{110} + a_{010} a_{100} a_{011} a_{101}) \\ &\quad + 4(a_{000} a_{011} a_{101} a_{110} + a_{001} a_{010} a_{100} a_{111}) \\ &= a_0^2 a_7^2 + a_1^2 a_6^2 + a_2^2 a_5^2 + a_3^2 a_4^2 \end{aligned} \tag{3.1}$$

$$\begin{aligned} &-2(a_0 a_1 a_6 a_7 + a_0 a_2 a_5 a_7 + a_0 a_4 a_3 a_7 + a_1 a_2 a_5 a_6 + a_1 a_3 a_4 a_6 + a_2 a_3 a_4 a_5) \\ &\quad + 4(a_0 a_3 a_5 a_6 + a_1 a_2 a_4 a_7) \end{aligned} \tag{3.2}$$

where we have made the binary conversion 0, 1, 2, 3, 4, 5, 6, 7 for 000, 001, 010, 011, 100, 101, 110, 111.

The hyperdeterminant vanishes iff the following system of equations in six unknowns p^A, q^B, r^D has a nontrivial solution, not allowing any of the pairs to be both zero:

$$\begin{aligned} a_{ABD} p^A q^B &= 0 \\ a_{ABD} p^A r^D &= 0 \\ a_{ABD} q^B r^D &= 0 \end{aligned} \tag{3.3}$$

For our purposes, the important properties of the hyperdeterminant are that it is a quartic invariant under $[SL(2)]^3$ and under a triality that interchanges A, B and D . These properties are valid whether the a_{ABD} are complex, real or integer.

One way to understand this triality is to think of having three different metrics (Alice, Bob and Daisy)

$$\begin{aligned} (\gamma_A)_{A_1 A_2} &= \epsilon^{B_1 B_2} \epsilon^{D_1 D_2} a_{A_1 B_1 D_1} a_{A_2 B_2 D_2} \\ (\gamma_B)_{B_1 B_2} &= \epsilon^{D_1 D_2} \epsilon^{A_1 A_2} a_{A_1 B_1 D_1} a_{A_2 B_2 D_2} \\ (\gamma_D)_{D_1 D_2} &= \epsilon^{A_1 A_2} \epsilon^{B_1 B_2} a_{A_1 B_1 D_1} a_{A_2 B_2 D_2} \end{aligned} \tag{3.4}$$

Explicitly,

$$\gamma = \begin{pmatrix} 2(a_0 a_6 - a_2 a_4) & a_0 a_7 - a_2 a_5 + a_1 a_6 - a_3 a_4 \\ a_0 a_7 - a_2 a_5 + a_1 a_6 - a_3 a_4 & 2(a_1 a_7 - a_3 a_5) \end{pmatrix} \tag{3.5}$$

$$\beta = \begin{pmatrix} 2(a_0a_3 - a_1a_2) & a_0a_7 - a_1a_6 + a_4a_3 - a_5a_2 \\ a_0a_7 - a_1a_6 + a_4a_3 - a_5a_2 & 2(a_4a_7 - a_5a_6) \end{pmatrix} \quad (3.6)$$

$$\alpha = \begin{pmatrix} 2(a_0a_5 - a_4a_1) & a_0a_7 - a_4a_3 + a_2a_5 - a_6a_1 \\ a_0a_7 - a_4a_3 + a_2a_5 - a_6a_1 & 2(a_2a_7 - a_6a_3) \end{pmatrix} \quad (3.7)$$

All are equivalent, however, since

$$\det \alpha = \det \beta = \det \gamma = -\text{Det } a \quad (3.8)$$

If we make the identifications

$$\begin{aligned} a_0 &= \frac{1}{\sqrt{2}}(-P^0 + P^2) \\ a_1 &= \frac{1}{\sqrt{2}}(-Q^0 + Q^2) \\ a_2 &= \frac{1}{\sqrt{2}}(P^1 - P^3) \\ a_3 &= \frac{1}{\sqrt{2}}(Q^1 - Q^3) \\ a_4 &= \frac{1}{\sqrt{2}}(-P^1 - P^3) \\ a_5 &= \frac{1}{\sqrt{2}}(-Q^1 - Q^3) \\ a_6 &= \frac{1}{\sqrt{2}}(-P^0 - P^2) \\ a_7 &= \frac{1}{\sqrt{2}}(-Q^0 - Q^2) \end{aligned} \quad (3.9)$$

then we find the $O(2,2)$ scalar products

$$\begin{aligned} 2(a_0a_6 - a_2a_4) &= (P^0)^2 + (P^1)^2 - (P^2)^2 - (P^3)^2 = P^2 \\ 2(a_1a_7 - a_3a_5) &= (Q_0)^2 + (Q_1)^2 - (Q_2)^2 - (Q_3)^2 = Q^2 \\ a_0a_7 - a_2a_5 + a_1a_6 - a_3a_4 &= (P^0Q_0) + (P^1Q_1) + (P^2Q_2) + (P^3Q_3) = P.Q \end{aligned}$$

so

$$\gamma = \begin{pmatrix} P^2 & P.Q \\ P.Q & Q^2 \end{pmatrix} \quad (3.10)$$

and

$$-\text{Det } a = P^2Q^2 - (P.Q)^2$$

4 Black hole entropy

The STU model admits extremal black hole solutions satisfying the Bogomolnyi mass formula. As usual, their entropy is given by one quarter the area of the event horizon. However, to calculate this area requires evaluating the mass not with the asymptotic values of the moduli, but with their frozen values on the horizon which are fixed in terms of the charges [27]. This ensures that the entropy is moduli-independent, as it should be. The relevant calculation was carried out in [11] for the model with the STU prepotential. The electric

and magnetic charges of that paper are denoted $(p^0, q_0), (p^1, q_1), (p^2, q_2), (p^3, q_3)$. In these variables, the entropy is given by

$$S = \pi \left(W(p^\Lambda, q_\Lambda) \right)^{1/2} \quad (4.1)$$

where

$$W(p^\Lambda, q_\Lambda) = -(p \cdot q)^2 + 4((p^1 q_1)(p^2 q_2) + (p^1 q_1)(p^3 q_3) + (p^3 q_3)(p^2 q_2)) - 4p^0 q_1 q_2 q_3 + 4q_0 p^1 p^2 p^3. \quad (4.2)$$

The function $W(p^\Lambda, q_\Lambda)$ is symmetric under transformations: $p^1 \leftrightarrow p^2 \leftrightarrow p^3$ and $q_1 \leftrightarrow q_2 \leftrightarrow q_3$. For the solution to be BPS we have to require $W > 0$.

If we make the identifications [1]

$$\begin{bmatrix} p^0 \\ p^1 \\ p^2 \\ p^3 \\ q_0 \\ q_1 \\ q_2 \\ q_3 \end{bmatrix} = \begin{bmatrix} -a_0 \\ -a_1 \\ -a_2 \\ a_4 \\ -a_7 \\ a_6 \\ a_5 \\ -a_3 \end{bmatrix} \quad (4.3)$$

we recognize from (3) that

$$W = -\text{Det } a \quad (4.4)$$

and hence the black hole entropy is given by

$$S = \pi \sqrt{-\text{Det } a} \quad (4.5)$$

Some examples of supersymmetric black hole solutions [29] are provided by the electric Kaluza-Klein black hole with $\alpha = (1, 0, 0, 0)$ and $\beta = (0, 0, 0, 0)$; the electric winding black hole with $\alpha = (0, 0, 0, -1)$ and $\beta = (0, 0, 0, 0)$; the magnetic Kaluza-Klein black hole with $\alpha = (0, 0, 0, 0)$ and $\beta = (0, -1, 0, 0)$; the magnetic winding black hole with $\alpha = (0, 0, 0, 0)$ and $\beta = (0, 0, -1, 0)$. These are characterized by a scalar-Maxwell coupling parameter $a = \sqrt{3}$. By combining these 1-particle states, we may build up 2-, 3- and 4-particle bound states at threshold [29, 9]. For example $\alpha = (1, 0, 0, -1)$ and $\beta = (0, 0, 0, 0)$ with $a = 1$; $\alpha = (1, 0, 0, -1)$ and $\beta = (0, -1, 0, 0)$ with $a = 1/\sqrt{3}$; $\alpha = (1, 0, 0, -1)$ and $\beta = (0, -1, -1, 0)$ with $a = 0$. The 1-, 2- and 3-particle states all yield vanishing contributions to $\text{Det } a$. A non-zero value is obtained for the 4-particle example, however, which is just the Reissner-Nordstrom black hole.

5 The N=8 generalization

The black holes described by Cayley's hyperdeterminant are those of $N = 2$ supergravity coupled to three vector multiplets, where the symmetry is $[SL(2, Z)]^3$. One might therefore ask whether the black hole/information theory correspondence could be generalized. There are three generalizations we might consider:

1) $N = 2$ supergravity coupled to l vector multiplets where the symmetry is $SL(2, Z) \times SO(l - 1, 2, Z)$ and the black holes carry charges belonging to the $(2, l + 1)$ representation ($l + 1$ electric plus $l + 1$ magnetic).

2) $N = 4$ supergravity coupled to m vector multiplets where the symmetry is $SL(2, Z) \times SO(6, 6 + m, Z)$ where the black holes carry charges belonging to the $(2, 12 + m)$ representation ($m + 12$ electric plus $m + 12$ magnetic).

3) $N = 8$ supergravity where the symmetry is the non-compact exceptional group $E_{7(7)}(Z)$ and the black holes carry charges belonging to the fundamental 56-dimensional representation (28 electric plus 28 magnetic).

In all three case there exit quartic invariants akin to Cayley's hyperdeterminant whose square root yields the corresponding black hole entropy. If there is to be a quantum information theoretic interpretation, however, it cannot just be random entanglement of more qubits, because the general n qubit entanglement is described by the group $[SL(2, C)]^n$, which, even after replacing Z by C , differs from the above symmetries (except when $n = 3$, which correspond to case (1) above with $l = 3$, the case we already know.).

We note, however, that

$$E_{7(7)}(Z) \supset [SL(2, Z)]^7 \quad (5.1)$$

and

$$E_7(C) \supset [SL(2, C)]^7, \quad (5.2)$$

We shall now show that the corresponding system in quantum information theory is that of seven qubits (Alice, Bob, Charlie, Daisy, Emma, Fred and George). However, the larger symmetry requires that they undergo at most tripartite entanglement of a very specific kind. The entanglement measure will be given by the quartic Cartan $E_7(C)$ invariant [16, 17, 18, 19].

6 Decomposition of $E_{7(7)}$

Consider the decomposition of the fundamental 56-dimensional representation of $E_{7(7)}$ under its maximal subgroup

$$\begin{aligned} E_{7(7)} &\supset SL(2)_A \times SO(6, 6) \\ 56 &\rightarrow (2, 12) + (1, 32) \end{aligned} \quad (6.1)$$

Further decomposing $SO(6, 6)$,

$$\begin{aligned} SL(2)_A \times SO(6, 6) &\supset SL(2)_A \times SL(2)_B \times SL(2)_D \times SO(4, 4) \\ (2, 12) + (1, 32) &\rightarrow (2, 2, 2, 1) \\ &+ (2, 1, 1, 8_v) + (1, 2, 1, 8_s) + (1, 1, 2, 8_c) \end{aligned} \quad (6.2)$$

Further decomposing $SO(4, 4)$,

$$\begin{aligned} SL(2)_A \times SL(2)_B \times SL(2)_D \times SO(4, 4) &\supset SL(2)_A \times SL(2)_B \times SL(2)_D \\ &\times SO(2, 2) \times SO(2, 2) \end{aligned}$$

$$\begin{aligned}
& (2, 2, 2, 1) + (2, 1, 1, 8_v) + (1, 2, 1, 8_s) + (1, 1, 2, 8_c) \rightarrow \\
& (2, 2, 2, 1, 1) + (2, 1, 1, 4, 1) + (2, 1, 1, 1, 4) \\
& + (1, 2, 1, 2, 2) + (1, 2, 1, 2, 2) + (1, 1, 2, 2, 2) + (1, 1, 2, 2, 2)
\end{aligned} \tag{6.3}$$

Finally, further decomposing each $SO(2, 2)$

$$\begin{aligned}
& SL(2)_A \times SL(2)_B \times SL(2)_D \times SO(2, 2) \times SO(2, 2) \supset \\
& SL(2)_A \times SL(2)_B \times SL(2)_D \times SL(2)_C \times SL(2)_G \times SL(2)_F \times SL(2)_E \\
& (2, 2, 2, 1, 1) + (2, 1, 1, 4, 1) + (2, 1, 1, 1, 4) \\
& + (1, 2, 1, 2, 2) + (1, 2, 1, 2, 2) + (1, 1, 2, 2, 2) + (1, 1, 2, 2, 2) \rightarrow \\
& (2, 2, 2, 1, 1, 1, 1) + (2, 1, 1, 2, 2, 1, 1) + (2, 1, 1, 1, 1, 2, 2) + \\
& (1, 2, 1, 2, 1, 1, 2) + (1, 2, 1, 1, 2, 2, 1) + (1, 1, 2, 2, 1, 2, 1) + (1, 1, 2, 1, 2, 1, 2)
\end{aligned}$$

In summary,

$$E_{7(7)} \supset SL(2)_A \times SL(2)_B \times SL(2)_C \times SL(2)_D \times SL(2)_E \times SL(2)_F \times SL(2)_G \tag{6.4}$$

and the 56 decomposes as

$$\begin{aligned}
& 56 \rightarrow \\
& (2, 2, 1, 2, 1, 1, 1) \\
& + (1, 2, 2, 1, 2, 1, 1) \\
& + (1, 1, 2, 2, 1, 2, 1) \\
& + (1, 1, 1, 2, 2, 1, 2) \\
& + (2, 1, 1, 1, 2, 2, 1) \\
& + (1, 2, 1, 1, 1, 2, 2) \\
& + (2, 1, 2, 1, 1, 1, 2)
\end{aligned} \tag{6.5}$$

An analogous decomposition holds for

$$E_7(C) \supset [SL(2, C)]^7. \tag{6.6}$$

7 Tripartite entanglement of 7 qubits

We have seen that in the case of three qubits, the tripartite entanglement is described by $[SL(2, C)]^3$ and that the entanglement measure is given by Cayley's hyperdeterminant. Now we consider seven qubits (Alice, Bob, Charlie, Daisy, Emma, Fred and George) but where Alice has tripartite entanglement not only with Bob/Daisy but also with Emma/Fred and also with George/Charlie, and similarly for the other six individuals. So, in fact, each person has tripartite entanglement with each of the remaining three couples:

$$\begin{aligned}
 |\Psi\rangle = & \\
 & a_{ABD}|ABD\rangle \\
 & +b_{BCE}|BCE\rangle \\
 & +c_{CDF}|CDF\rangle \\
 & +d_{DEG}|DEG\rangle \\
 & +e_{EFA}|EFA\rangle \\
 & +f_{FGB}|FGB\rangle \\
 & +g_{GAC}|GAC\rangle
 \end{aligned} \tag{7.1}$$

Note that:

- 1) Any pair of states has an individual in common
- 2) Each individual is excluded from four out of the seven states
- 3) Two given individuals are excluded from two out of the seven states
- 4) Three given individuals are never excluded

The entanglement may be represented by a heptagon with vertices A,B,C,D,E,F,G and seven triangles ABD, BCE, CDF, DEG, EFA, FGB, and GAC. See Figure 1. Alternatively, we can use the Fano plane. See Figure 2. The Fano plane corresponds to the multiplication table of the octonions as may be seen from the description of the state $|\Psi\rangle$ given in Table 1.

Each of the seven states transforms as a $(2, 2, 2)$ under three of the $SL(2)$'s and are singlets under the remaining four. Note that from (6.2) we see that the A-B-C triality of section 3 is linked with the $8_v - 8_s - 8_c$ triality of the $SO(4, 4)$. For example, interchanging A and B leaves $|\Psi\rangle$ invariant provided we also interchange C and F. Individually, therefore, the tripartite entanglement of each of the seven states is given by Cayley's hyperdeterminant. Taken together however, we see from (6.5) that they transform as a complex 56 of $E_7(C)$. Their tripartite entanglement must be is given by an expression that is quartic in the coefficients a, b, c, d, e, f, g and invariant under $E_7(C)$. The unique possibility is the Cartan invariant I_4 , and so the 3-tangle is given by

$$\tau_3(ABCDEFG) = 4|I_4| \tag{7.2}$$

If the wave-function (7.1) is normalized, then $0 \leq \tau_3(ABCDEFG) \leq 1$.




Figure 1: The E_7 entanglement diagram. Each of the seven vertices A,B,C,D,E,F,G represents a qubit and each of the seven triangles ABD, BCE, CDF, DEG, EFA, FGB, GAC describes a tripartite entanglement.

8 Cartan's $E_{7(7)}$ invariant

The Cremmer-Julia [17] form of the Cartan $E_{7(7)}$ invariant may be written as

$$I_4 = \text{Tr}(Z\bar{Z})^2 - \frac{1}{4}(\text{Tr } Z\bar{Z})^2 + 4(\text{Pf } Z + \text{Pf } \bar{Z}) , \quad (8.1)$$

and the Cartan form [16] may be written as

$$I_4 = -\text{Tr}(x y)^2 + \frac{1}{4}(\text{Tr } x y)^2 - 4(\text{Pf } x + \text{Pf } y) . \quad (8.2)$$

Here

$$Z_{AB} = -\frac{1}{4\sqrt{2}}(x^{ab} + iy_{ab})(\Gamma^{ab})_{AB} \quad (8.3)$$

and

$$x^{ab} + iy_{ab} = -\frac{\sqrt{2}}{4}Z_{AB}(\Gamma^{AB})_{ab} \quad (8.4)$$

The matrices of the $SO(8)$ algebra are $(\Gamma^{ab})_{AB}$ where $(a b)$ are the 8 vector indices and (A, B) are the 8 spinor indices. The $(\Gamma^{ab})_{AB}$ matrices can be considered also as $(\Gamma^{AB})_{ab}$ matrices due to equivalence of the vector and spinor representations of the $SO(8)$ Lie algebra. The exact relation between the Cartan invariant in (8.2) and Cremmer-Julia invariant [17] in (8.1) was established in [20, 21]. The quartic invariant I_4 of $E_{7(7)}$ is also related to the octonionic Jordan algebra J_3^O [19].

In the stringy black hole context, Z_{AB} is the central charge matrix and (x, y) are the quantized charges of the black hole (28 electric and 28 magnetic). The relation between the entropy of stringy black holes and the Cartan-Cremmer-Julia $E_{7(7)}$ invariant was established




Figure 2: The Fano plane has seven points, representing the seven qubits, and seven lines (the circle counts as a line) with three points on every line, representing the tripartite entanglement, and three lines through every point.

in [18]. The central charge matrix Z_{AB} can be brought to the canonical basis for the skew-symmetric matrix using an $SU(8)$ transformation:

$$Z_{ab} = \begin{pmatrix} z_1 & 0 & 0 & 0 \\ 0 & z_2 & 0 & 0 \\ 0 & 0 & z_3 & 0 \\ 0 & 0 & 0 & z_4 \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad (8.5)$$

where $z_i = \rho_i e^{i\varphi_i}$ are complex. In this way the number of entries is reduced from 56 to 8. In a systematic treatment in [22], the meaning of these parameters was clarified. From 4 complex values of $z_i = \rho_i e^{i\varphi_i}$ one can remove 3 phases by an $SU(8)$ rotation, but the overall phase cannot be removed; it is related to an extra parameter in the class of black hole solutions [23, 24]. In this basis, the quartic invariant takes the form [18]

$$\begin{aligned} I_4 &= \sum_i |z_i|^4 - 2 \sum_{i < j} |z_i|^2 |z_j|^2 + 4(z_1 z_2 z_3 z_4 + \bar{z}_1 \bar{z}_2 \bar{z}_3 \bar{z}_4) \\ &= (\rho_1 + \rho_2 + \rho_3 + \rho_4)(\rho_1 + \rho_2 - \rho_3 - \rho_4)(\rho_1 - \rho_2 + \rho_3 - \rho_4)(\rho_1 - \rho_2 - \rho_3 + \rho_4) \\ &\quad + 8\rho_1 \rho_2 \rho_3 \rho_4 (\cos \varphi - 1) \end{aligned} \quad (8.6)$$

Therefore a 5-parameter solution is called a generating solution for other black holes in N=8 supergravity/M-theory. The expression for their entropy is always given by

$$S = \pi \sqrt{|I_4|} \quad (8.7)$$

for some subset of 5 of the 8 parameters mentioned above. Recently a new class of solutions was discovered, describing black rings. The maximal number of parameters for the known

	A	B	C	D	E	F	G
A		D	G	-B	F	-E	-C
B	-D		E	A	-C	G	-F
C	-G	-E		F	B	-D	A
D	B	-A	-F		G	C	-E
E	-F	C	-B	-G		A	D
F	E	-G	D	-C	-A		B
G	C	F	-A	E	-D	-B	

Table 1: The entanglement of the state $|\Psi\rangle$ coincides with the multiplication table of the octonions.

solutions is 7. The entropy of black ring solutions found so far was identified in [25, 26] with the expression (8.7) for a subset of 7 out of 8 parameters mentioned above.

Kallosh and Linde have shown that I_4 depending on 4 complex eigenvalues can be represented as Cayley's hyperdeterminant of a hypermatrix a_{ABD} . To see this, we that in x, y basis only the $SO(8)$ symmetry is manifest, which means that every term in (8.2) is invariant only under $SO(8)$ symmetry. However, it was proved in [16] and [17] that the sum of all terms in (8.2) is invariant under the full $SU(8)$ symmetry, which acts as follows

$$\delta(x^{ab} \pm iy_{ab}) = (2\Lambda^a_{[c}\delta^b]_{d]} \pm i\Sigma_{abcd})(x^{cd} \mp iy_{cd}) . \quad (8.8)$$

The total number of parameters is 63, where 28 are from the manifest $SO(8)$ and 35 from the antisymmetric self-dual $\Sigma_{abcd} = *\Sigma^{abcd}$. Thus one can use the $SU(8)$ transformation of the complex matrix $x^{ab} + iy_{ab}$ and bring it to the canonical form with some complex eigenvalues $\lambda_I, I = 1, 2, 3, 4$. The value of the quartic invariant (8.2) will not change.

$$(x^{ab} + iy_{ab})_{\text{can}} = \begin{pmatrix} 0 & \lambda_1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -\lambda_1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \lambda_2 & 0 & 0 & 0 & 0 \\ 0 & 0 & -\lambda_2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & \lambda_3 & 0 & 0 \\ 0 & 0 & 0 & 0 & -\lambda_3 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & \lambda_4 \\ 0 & 0 & 0 & 0 & 0 & 0 & -\lambda_4 & 0 \end{pmatrix} \quad (8.9)$$

The relation between the complex coefficients λ_I , the parameters x^{ij} and y_{kl} , the matrix a_{ABD} and the black hole charges p^i and q_k [1] is given by the following dictionary:

$$\lambda_1 = x^{12} + iy_{12} = a_{111} + ia_{000} = -q_0 - ip^0$$

$$\begin{aligned}
\lambda_2 &= x^{34} + iy_{34} = a_{001} + ia_{110} = -p^1 + iq_1 \\
\lambda_3 &= x^{56} + iy_{56} = a_{010} + ia_{101} = -p^2 + iq_2 \\
\lambda_4 &= x^{78} + iy_{78} = a_{100} + ia_{011} = p^3 - iq_3
\end{aligned} \tag{8.10}$$

If we now write the quartic $E_{7(7)}$ Cartan invariant in the canonical basis (x^{ij}, y_{ij}) , $i, j = 1, \dots, 8$:

$$\begin{aligned}
I_4 &= -(x^{12}y_{12} + x^{34}y_{34} + x^{56}y_{56} + x^{78}y_{78})^2 - 4(x^{12}x^{34}x^{56}x^{78} + y_{12}y_{34}y_{56}y_{78}) \\
&\quad + 4(x^{12}x^{34}y_{12}y_{34} + x^{12}x^{56}y_{12}y_{56} + x^{34}x^{56}y_{34}y_{56} + x^{12}x^{78}y_{12}y_{78} + x^{34}x^{78}y_{34}y_{78} \\
&\quad + x^{56}x^{78}y_{56}y_{78}) .
\end{aligned} \tag{8.11}$$

then it may now be compared to Cayley's hyperdeterminant (3.1). We find

$$I_4 = -\text{Det } a \tag{8.12}$$

The above discussion of $E_{7(7)}$ also applies, mutatis mutandis, to $E_7(C)$.

To understand better the entanglement we note that, as a result of (6.5), Cartan's invariant contains not one Cayley hyperdeterminant but seven! It may be written as the sum of seven terms each of which is invariant under $[SL(2)]^3$ plus cross terms. To see this, denote a 2 in one of the seven entries in (6.5) by A, B, C, D, E, F, G. So we may rewrite (6.5) as

$$56 = (ABD) + (BCE) + (CDF) + (DEG) + (EFA) + (FGB) + (GAC) \tag{8.13}$$

or symbolically

$$56 = a + b + c + d + e + f + g \tag{8.14}$$

Then I_4 is the singlet in $56 \times 56 \times 56 \times 56$:

$$\begin{aligned}
J_4 &\sim a^4 + b^4 + c^4 + d^4 + e^4 + f^4 + g^4 + \\
&\quad 2[a^2b^2 + b^2c^2 + c^2d^2 + d^2e^2 + e^2f^2 + f^2g^2 + g^2a^2 + \\
&\quad a^2c^2 + b^2d^2 + c^2e^2 + d^2f^2 + e^2g^2 + f^2a^2 + g^2b^2 + \\
&\quad a^2d^2 + b^2e^2 + c^2f^2 + d^2g^2 + e^2a^2 + f^2b^2 + g^2c^2] \\
&\quad + 8[bcdf + cdeg + defa + efgb + fgac + gabd + abce]
\end{aligned} \tag{8.15}$$

where products like

$$\begin{aligned}
a^4 &= (ABD)(ABD)(ABD)(ABD) \\
&= \epsilon^{A_1A_2}\epsilon^{B_1B_2}\epsilon^{D_1D_4}\epsilon^{A_3A_4}\epsilon^{B_3B_4}\epsilon^{D_2D_3}a_{A_1B_1D_1}a_{A_2B_2D_2}a_{A_3B_3D_3}a_{A_4B_4D_4}
\end{aligned} \tag{8.16}$$

exclude four individuals (here Charlie, Emma, Fred and George), products like

$$\begin{aligned}
a^2f^2 &= (ABD)(ABD)(FGB)(FGB) \\
&= \epsilon^{A_1A_2}\epsilon^{B_1B_2}\epsilon^{D_1D_4}\epsilon^{F_3F_4}\epsilon^{G_3G_4}\epsilon^{D_2B_3}a_{A_1B_1D_1}a_{A_2B_2D_2}f_{F_3G_3B_3}f_{F_4G_4B_4}
\end{aligned} \tag{8.17}$$

exclude two individuals (here Charlie and Emma), and products like

$$\begin{aligned}
abce &= (ABD)(BCE)(CDF)(EFA) \\
&= \epsilon^{A_1A_4}\epsilon^{B_1B_2}\epsilon^{C_2C_3}\epsilon^{D_1D_3}\epsilon^{E_2E_4}\epsilon^{F_3F_4}a_{A_1B_1D_1}b_{B_2C_2E_2}c_{C_3D_3F_3}e_{E_4F_4A_4}
\end{aligned} \tag{8.18}$$

exclude one individual (here George).

9 The black hole analogy

In the STU stringy black hole context [1, 9, 11, 2] the a_{ABC} are integers (corresponding to quantized charges) and hence the symmetry group is $[SL(2, Z)]^3$ rather than $[SL(2, C)]^3$. However, as discussed by Levay [3], there is a branch of quantum information theory which concerns itself with real qubits, called *rebits*, for which the a_{ABC} are real. (One difference remains, however: one may normalize the wave function, whereas for black holes there is no such restriction on the charges a_{ABC} .) It turns out that there are three reality classes which can be characterized by the hyperdeterminant

$$\begin{aligned} 1) & \text{ Det } a < 0 \\ 2) & \text{ Det } a = 0 \\ 3) & \text{ Det } a > 0 \end{aligned} \tag{9.1}$$

Case (1) corresponds to the non-separable or GHZ class [28], for example,

$$|\Psi\rangle = \frac{1}{2}(-|000\rangle + |011\rangle + |101\rangle + |110\rangle) \tag{9.2}$$

Case (2) corresponds to the separable (A-B-C, A-BC, B-CA, C-AB) and W classes, for example

$$|\Psi\rangle = \frac{1}{\sqrt{3}}(|100\rangle + |010\rangle + |001\rangle) \tag{9.3}$$

In the string/supergravity interpretation [1], cases (1) and (2) were shown to correspond to BPS black holes, for which half of the supersymmetry is preserved. Case (1) has non-zero horizon area and entropy (“large” black holes), and case (2) to vanishing horizon area and entropy (“small” black holes), at least at the semi-classical level. However, small black holes may acquire a non-zero entropy through higher order quantum effects. This entropy also has a quantum information interpretation involving bipartite entanglement of the three qubits [2].

Case (3) is also GHZ, for example the above GHZ state (9.2) with a sign flip

$$|\Psi\rangle = \frac{1}{2}(|000\rangle + |011\rangle + |101\rangle + |110\rangle) \tag{9.4}$$

In the string/supergravity interpretation, case (3) corresponds to non-BPS black holes [2]. With four non-zero charges (q_0, p^1, p^2, p^3) in (8.10), for example, an extreme but non-BPS black hole [29] may be obtained by flipping the sign [30] of one of the charges. The canonical GHZ state

$$|\Psi\rangle = \frac{1}{\sqrt{2}}|111\rangle + \frac{1}{\sqrt{2}}|000\rangle \tag{9.5}$$

also belongs to case (3).

In the $N = 8$ theory, “large” and “small” black holes are classified by the sign of I_4 :

$$\begin{aligned} 1) & I_4 > 0 \\ 2) & I_4 = 0 \end{aligned}$$

$$3) I_4 < 0 \quad (9.6)$$

Once again, non-zero I_4 corresponds to large black holes, which are BPS for $I_4 > 0$ and non-BPS for $I_4 < 0$, and vanishing I_4 to small black holes. However, in contrast to $N = 2$, case (1) requires that only 1/8 of the supersymmetry is preserved, while we may have 1/8, 1/4 or 1/2 for case (2).

It is worth noting that the charge orbits corresponding to non-zero I_4 are associated with the following cosets:

$$\frac{E_{7(7)}}{E_{6(2)}} \quad (9.7)$$

and

$$\frac{E_{7(7)}}{E_{6(6)}} \quad (9.8)$$

The large black hole solutions can be found [31] by solving the $N = 8$ classical attractor equations [27] when at the attractor value the Z_{AB} matrix, in normal form, becomes

$$Z_{AB} = \begin{pmatrix} Z\epsilon & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (9.9)$$

for positive I_4 and

$$Z_{AB} = e^{i\pi/4} |Z| \begin{pmatrix} \epsilon & 0 & 0 & 0 \\ 0 & \epsilon & 0 & 0 \\ 0 & 0 & \epsilon & 0 \\ 0 & 0 & 0 & \epsilon \end{pmatrix} \quad (9.10)$$

for negative I_4 . These values exhibit the maximal compact symmetries $SU(6) \times SU(2)$ and $USp(8)$ for the positive and negative I_4 , respectively.

If the phase in (8.6) vanishes (which is the case if the configuration preserves at least 1/4 supersymmetry [22]), I_4 becomes

$$I_4 = \lambda_1 \lambda_2 \lambda_3 \lambda_4, \quad (9.11)$$

where we have defined λ_i by

$$\begin{aligned} \lambda_1 &= \rho_1 + \rho_2 + \rho_3 + \rho_4 \\ \lambda_2 &= \rho_1 + \rho_2 - \rho_3 - \rho_4 \\ \lambda_3 &= \rho_1 - \rho_2 + \rho_3 - \rho_4 \\ \lambda_4 &= \rho_1 - \rho_2 - \rho_3 + \rho_4 \end{aligned} \quad (9.12)$$

and we order the λ_i so that $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq |\lambda_4|$. The charge orbits for the small black holes depend on the number of unbroken supersymmetries or the number of vanishing eigenvalues. The orbit is [19, 22, 32]

$$\frac{E_{7(7)}}{H_{1,2,3}} \quad (9.13)$$

where

$$H_1 = F_{4(4)} \ltimes T_{26} \quad \lambda_1, \lambda_2, \lambda_3 \neq 0, \quad \lambda_4 = 0 \quad (1/8 \text{ BPS})$$

$$\begin{aligned}
H_2 &= SO(5, 6) \ltimes (T_{32} \times T_1) & \lambda_1, \lambda_2 \neq 0, \quad \lambda_3, \lambda_4 = 0 & \quad (1/4 \text{ BPS}) \\
H_3 &= E_{6(6)} \ltimes T_{27} & \lambda_1 \neq 0, \quad \lambda_2, \lambda_3, \lambda_4 = 0 & \quad (1/2 \text{ BPS})
\end{aligned} \tag{9.14}$$

For $N = 8$, as for $N = 2$, the large black holes correspond to the two classes of GHZ-type (entangled) states and small black holes to the separable or W class.

10 Subsectors

Having understood the analogy between $N = 8$ black holes and the tripartite entanglement of 7 qubits using $E_{7(7)}$, we may now find the analogy in the $N = 4$ case using $SL(2) \times SO(6, 6)$ and the $N = 2$ case using $SL(2) \times SO(2, 2)$.

For $N = 4$, as may be seen from (6.2), we still have an $[SL(2)]^7$ subgroup but now there are only 24 states

$$|\Psi\rangle = a_{ABD}|ABD\rangle + e_{EFA}|EFA\rangle + g_{GAC}|GAC\rangle \tag{10.1}$$

So only Alice talks to all the others. This is described by just those three lines passing through A in the Fano plane. Then the equations analogous to (8.13) and (8.14) are

$$(2, 12) = (ABD) + (EFA) + (GAC) = a + e + g \tag{10.2}$$

and the corresponding quartic invariant, I_4 , reduces to the singlet in $(2, 12) \times (2, 12) \times (2, 12) \times (2, 12)$.

$$I_4 \sim a^4 + e^4 + g^4 + 2[e^2g^2 + g^2a^2 + a^2e^2] \tag{10.3}$$

If we identify the 24 numbers $(a_{ABD}, e_{EFA}, g_{GAC})$ with (P^μ, Q_ν) with $\mu, \nu = 0, \dots, 11$, this becomes [9, 23, 24]

$$I_4 = P^2Q^2 - (P \cdot Q)^2 \tag{10.4}$$

which is manifestly invariant under $SL(2) \times SO(6, 6)$.

For $N = 2$, as may be seen from (6.2), we only have an $[SL(2)]^3$ subgroup and there are only 8 states

$$|\Psi\rangle = a_{ABD}|ABD\rangle \tag{10.5}$$

This is described by just the ABD line in the Fano plane. This is simply the usual tripartite entanglement, for which

$$(2, 2, 2) = (ABD) = a \tag{10.6}$$

and the corresponding quartic invariant

$$I_4 \sim a^4 \tag{10.7}$$

is just Cayley's hyperdeterminant

$$I_4 = -\text{Deta} \tag{10.8}$$

11 Conclusions

We note that the 56-dimensional Hilbert space given in (6.5) and (7.1) is not a subspace of the usual 2^7 -dimensional seven-qubit Hilbert space given by $(2, 2, 2, 2, 2, 2, 2)$, but rather a direct sum of seven 2^3 -dimensional three-qubit Hilbert spaces $(2, 2, 2)$. This is however, a subspace of the 3^7 -dimensional seven-qutrit Hilbert space given by $(3, 3, 3, 3, 3, 3, 3)$. Under

$$[SL(3)]^7 \rightarrow [SL(2)]^7 \quad (11.1)$$

we have the decomposition

$$\begin{aligned} (3, 3, 3, 3, 3, 3, 3) \rightarrow \\ & 1 \text{ term like } (2, 2, 2, 2, 2, 2, 2) \\ & 7 \text{ terms like } (2, 2, 2, 2, 2, 2, 1) \\ & 21 \text{ terms like } (2, 2, 2, 2, 2, 1, 1) \\ & 35 \text{ terms like } (2, 2, 2, 2, 1, 1, 1) \\ & 35 \text{ terms like } (2, 2, 2, 1, 1, 1, 1) \\ & 21 \text{ terms like } (2, 2, 1, 1, 1, 1, 1) \\ & 7 \text{ terms like } (2, 1, 1, 1, 1, 1, 1) \\ & 1 \text{ term like } (1, 1, 1, 1, 1, 1, 1) \end{aligned} \quad (11.2)$$

which contains

$$\begin{aligned} & (2, 2, 1, 2, 1, 1, 1) \\ & +(1, 2, 2, 1, 2, 1, 1) \\ & +(1, 1, 1, 2, 2, 1, 2) \\ & +(2, 1, 1, 1, 2, 2, 1) \\ & +(1, 2, 1, 1, 1, 2, 2) \\ & +(2, 1, 2, 1, 1, 1, 2) \end{aligned} \quad (11.3)$$

So the Fano plane entanglement we have described fits within conventional quantum information theory.

The Fano plane also finds application in switching networks that can connect any phone to any other phone. It is the 3-switching network for 7 numbers. However there also exists a 4-switching network for 13 numbers, a 5-switching network for 21 numbers, and generally an $(n+1)$ -switching network for (n^2+n+1) numbers corresponding to the projective planes of order n [33]. It would be worthwhile pursuing the corresponding quantum bit entanglements.

Exceptional groups, such as $E_{7(7)}$, have featured in supergravity, string theory, M-theory and other speculative attempts at unification of the fundamental forces. However, it is unusual to find an exceptional group appearing in the context of qubit entanglement. It would be interesting to see whether it can be subject to experimental test.

12 Acknowledgement

We are grateful to Murat Gunaydin for pointing out the connection between our entanglement diagram, Figure 1, and the multiplication table of the octonions. MJD has enjoyed useful conversations with Leron Borsten. Help with the diagrams from Laura Andrianopoli is also gratefully acknowledged.

This work was supported in part by the National Science Foundation under grant number PHY-0245337 and PHY-0555605. Any opinions, findings and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation. The work of S.F. has been supported in part by the European Community Human Potential Program under contract MRTN-CT-2004-005104 Constituents, fundamental forces and symmetries of the universe, in association with INFN Frascati National Laboratories and by the D.O.E grant DE-FG03-91ER40662, Task C. The work of MJD is supported in part by PPARC under rolling grant PPA/G/O/2002/00474, PP/D50744X/1.

References

- [1] M. J. Duff, “String triality, black hole entropy and Cayley’s hyperdeterminant,” arXiv:hep-th/0601134.
- [2] R. Kallosh and A. Linde, “Strings, black holes, and quantum information,” arXiv:hep-th/0602061.
- [3] P. Levay, “Stringy black holes and the geometry of entanglement,” Phys. Rev. D **74**, 024030 (2006) [arXiv:hep-th/0603136].
- [4] M. J. Duff and S. Ferrara, “ E_7 and the tripartite entanglement of seven qubits,” arXiv:quant-ph/0609227.
- [5] P. Levay, “Strings, black holes, the tripartite entanglement of seven qubits and the Fano plane,” arXiv:hep-th/0610314.
- [6] M. J. Duff and R. R. Khuri, “Four-Dimensional String / String Duality,” Nucl. Phys. B **411**, 473 (1994) [arXiv:hep-th/9305142].
- [7] C. M. Hull and P. K. Townsend, “Unity of superstring dualities,” Nucl. Phys. B **438**, 109 (1995) [arXiv:hep-th/9410167].
- [8] M. J. Duff, “Strong / weak coupling duality from the dual string,” Nucl. Phys. B **442**, 47 (1995) [arXiv:hep-th/9501030].
- [9] M. J. Duff, J. T. Liu and J. Rahmfeld, “Four-dimensional string-string-string triality,” Nucl. Phys. B **459**, 125 (1996) [arXiv:hep-th/9508094].
- [10] A. Cayley, “On the theory of linear transformations,” Camb. Math. J. 4 193-209, 1845.

- [11] K. Behrndt, R. Kallosh, J. Rahmfeld, M. Shmakova and W. K. Wong, “STU black holes and string triality,” *Phys. Rev. D* **54**, 6293 (1996) [arXiv:hep-th/9608059].
- [12] V. Coffman, J. Kundu and W. K. Wootters, “Distributed Entanglement,” *Phys. Rev. A* **61**, 052306 (2000) [arXiv:quant-ph/9907047].
- [13] A. Miyake and M. Wadati, “Multipartite entanglement and hyperdeterminants,” [arXiv:quant-ph/0212146].
- [14] M. J. Duff, “Hidden symmetries of the Nambu-Goto action,” arXiv:hep-th/0602160.
- [15] F. Klein, “Zur Theorie der Liniencomplexe des ersten und zweiten Grades”, *Math. Ann.* **2**, 198-226, 1870.
- [16] E. Cartan, “Oeuvres completes”, (Editions du Centre National de la Recherche Scientifique, Paris, 1984).
- [17] E. Cremmer and B. Julia, “The SO(8) Supergravity,” *Nucl. Phys. B* **159**, 141 (1979).
- [18] R. Kallosh and B. Kol, “E(7) Symmetric Area of the Black Hole Horizon,” *Phys. Rev. D* **53**, 5344 (1996) [arXiv:hep-th/9602014].
- [19] S. Ferrara and M. Gunaydin, “Orbits of exceptional groups, duality and BPS states in string theory,” *Int. J. Mod. Phys. A* **13**, 2075 (1998) [arXiv:hep-th/9708025].
- [20] V. Balasubramanian, F. Larsen and R. G. Leigh, “Branes at angles and black holes,” *Phys. Rev. D* **57**, 3509 (1998) [arXiv:hep-th/9704143];
- [21] M. Gunaydin, K. Koepsell and H. Nicolai, “Conformal and quasiconformal realizations of exceptional Lie groups,” *Commun. Math. Phys.* **221**, 57 (2001) [arXiv:hep-th/0008063].
- [22] S. Ferrara and J. M. Maldacena, “Branes, central charges and U -duality invariant BPS conditions,” *Class. Quant. Grav.* **15**, 749 (1998) [arXiv:hep-th/9706097].
- [23] M. Cvetič and D. Youm, “All the Static Spherically Symmetric Black Holes of Heterotic String on a Six Torus,” *Nucl. Phys. B* **472**, 249 (1996) [arXiv:hep-th/9512127].
- [24] M. Cvetič and A. A. Tseytlin, “Solitonic strings and BPS saturated dyonic black holes,” *Phys. Rev. D* **53**, 5619 (1996) [Erratum-ibid. *D* **55**, 3907 (1997)] [arXiv:hep-th/9512031].
- [25] I. Bena and P. Kraus, “Microscopic description of black rings in AdS/CFT,” *JHEP* **0412**, 070 (2004) [arXiv:hep-th/0408186].
- [26] I. Bena, P. Kraus and N. P. Warner, “Black rings in Taub-NUT,” *Phys. Rev. D* **72**, 084019 (2005) [arXiv:hep-th/0504142].
- [27] S. Ferrara, R. Kallosh and A. Strominger, “N=2 extremal black holes,” *Phys. Rev. D* **52**, 5412 (1995) [arXiv:hep-th/9508072].

- [28] D. M. Greenberger, M. Horne and A. Zeilinger, in *Bell's Theorem, Quantum Theory and Conceptions of the Universe*, ed. M. Kafatos (Kluwer, Dordrecht, 1989)
- [29] M. J. Duff and J. Rahmfeld, "Massive string states as extreme black holes," Phys. Lett. B **345**, 441 (1995) [arXiv:hep-th/9406105].
- [30] M. J. Duff and J. Rahmfeld, "Bound States of Black Holes and Other P-branes," Nucl. Phys. B **481**, 332 (1996) [arXiv:hep-th/9605085].
- [31] S. Ferrara and R. Kallosh, "On $N = 8$ attractors," Phys. Rev. D **73**, 125005 (2006) [arXiv:hep-th/0603247].
- [32] H. Lu, C. N. Pope and K. S. Stelle, "Multiplet structures of BPS solitons," Class. Quant. Grav. **15**, 537 (1998) [arXiv:hep-th/9708109].
- [33] E. Pegg Jr, "The Fano Plane", MAA online,
http://www.maa.org/editorial/mathgames/mathgames_05_30_06.html
- [34] A. Ceresole, R. D'Auria, S. Ferrara and A. Van Proeyen, "On electromagnetic duality in locally supersymmetric $N=2$ Yang-Mills theory," arXiv:hep-th/9412200.